



**ILLINOIS NATURAL
HISTORY SURVEY**
PRAIRIE RESEARCH INSTITUTE

Freshwater Mussels of the Mississippi South/ South Central Basins

Diane K. Shasteen, Alison L. Price, Sarah A. Bales

INHS Technical Report 2012 (08)

Prepared for:

Illinois Department of Natural Resources: Office of Resource Conservation
U.S. Fish & Wildlife Service
Illinois Natural History Survey

Issued March 7, 2012

Prairie Research Institute, University of Illinois at Urbana Champaign
William Shilts, Executive Director

Illinois Natural History Survey
Brian D. Anderson, Director
1816 South Oak Street
Champaign, IL 61820
217-333-6830



Freshwater Mussels of the Mississippi South/South Central Basins



2012

Illinois Natural History Survey, Prairie Research Institute, University of Illinois
Illinois Department of Natural Resources

Diane Shasteen, Alison Price, Sarah Bales

Preface

While broad geographic information is available on the distribution and abundance of mussels in Illinois, systematically collected mussel-community data sets required to integrate mussels into aquatic community assessments do not exist. In 2009, a project funded by a US Fish and Wildlife Service State Wildlife Grant was undertaken to survey and assess the freshwater mussel populations at wadeable sites from 33 stream basins in conjunction with the Illinois Department of Natural Resources (IDNR)/Illinois Environmental Protection Agency (IEPA) basin surveys. Inclusion of mussels into these basin surveys contributes to the comprehensive basin monitoring programs that include water and sediment chemistry, instream habitat, macroinvertebrate, and fish, which reflect a broad spectrum of abiotic and biotic stream resources. These mussel surveys will provide reliable and repeatable techniques for assessing the freshwater mussel community in sampled streams. These surveys also provide data for future monitoring of freshwater mussel populations on a local, regional, and watershed basis.

Agency Contacts

Kevin S. Cummings, INHS, ksc@inhs.illinois.edu (217) 333-1623
Bob Szafoni, IDNR, Robert.szafoni@illinois.gov, (217) 348-0175
Ann Marie Holtrop, IDNR, ann.holtrop@illinois.gov, (217) 785-4325

Suggested Citation

Shasteen, D.K., A.L. Price, and S.A. Bales. 2012. Freshwater Mussels of the Mississippi South/South Central River Basins. Illinois Natural History Survey Technical Report 2012 (08). Champaign, Illinois. 14 pp + appendix.

Acknowledgements

This study was supported by funding from the US Fish and Wildlife Service, State Wildlife Grant (T-53-D-1, Investigating Mussel Communities in Illinois Streams), the IDNR, and the Illinois Natural History Survey (INHS). Our extreme gratitude goes to the primary investigators for the project: Ann Holtrop, Kevin Cummings, Robert Szafoni, and Dr. Yong Cao, who served as our mentors and made this project possible. We would like to thank all people involved in our surveys, especially our field assistants (Rachel Vinsel, Jen Schwab, Hunter Ray, Andrew Berger, Andrew Repp, Jeff Gersch, Otis Foster and Colleen McLean), IDNR fisheries biologists, IEPA water monitoring biologists, and volunteers from other agencies. We would like to extend gratitude to all the landowners, both public and private, who allowed us access to their properties. We would like to thank Andrew Hulin for the creation of maps for this report and Christine Mayer for INHS Collection database support.

Introduction

Freshwater mussel populations have been declining for decades and are among the most seriously impacted aquatic animals worldwide (Bogan 1993, Williams et al. 1993). It is estimated that nearly 70% of the approximately 300 North American mussel taxa are extinct, federally-listed as endangered or threatened, or in need of conservation status (Williams et al. 1993, Strayer et al. 2004). In Illinois, 25 of the 62 extant species (44%) are listed as threatened or endangered (Illinois Endangered Species Protection Board 2011). While broad geographic information is available on the distribution and abundance of mussels in Illinois, systematically collected mussel-community data sets required to integrate mussels into aquatic community assessments do not exist. Sampling of mussels has been very sporadic and limited in the Mississippi South and South Central basins, and no known reports have been compiled on the mussel communities in these regions. This report summarizes the mussel survey conducted in the Mississippi South (MS) and Mississippi South Central (MSC) basins in 2009 and 2010 in conjunction with IDNR and IEPA basin surveys.

The Mississippi South basin consists of two separate drainages. The Mary's River drains an area of approximately 600 km² between the Kaskaskia River and the Big Muddy River and empties directly into the Mississippi River near Chester. To the south of the Big Muddy is the Clear Creek drainage, which drains approximately 1862 km² (Page et al. 1992). These two drainages occur in predominately rural areas with the major land use being agricultural (Page et al. 1992). The MS basin flows through three natural divisions including the Lower Mississippi River Bottomlands, Coastal Plains, and Ozark Southern divisions (Schwegman 1973). The Mississippi South Central basin includes the Cahokia Creek and Wood River drainages. The MSC basin, which flows through the area known as the American Bottoms, is highly impacted by urbanization, industry, and abandoned coal mines. The MSC basin flows through four natural divisions including the Western Forest-Prairie Division, Southern Till Plain, Middle Mississippi Border, and the Lower Mississippi River Bottomlands (Schwegman 1973). The MSC and MS basins combined drain the area from Honey Point Township in Macoupin County (headwaters of Cahokia Creek) to Thebes (Sammons Creek) in the southernmost county, Alexander, on the Mississippi.

Land-use and Instream Habitat

The primary land use of the Mary's River drainage is row-crop agriculture (75%) with 12% of the land being forested (IDA 2000). The city of Chester has a current population of approximately 8500 people, and is the only municipality located in the basin (US Census Bureau 2010). The Clear Creek drainage encompasses Union and Alexander counties in southwestern portion of the state. Land use in these counties includes approximately 30% forested lands, slightly over 50% agricultural and less than 2% urban (IDA 2000). In contrast, the MSC basin is located in the

Metro East region of St. Louis, which comprises the eastern suburbs of St. Louis, Missouri. It encompasses five southern Illinois counties including Clinton, Jersey, Madison, Monroe and St. Clair and has a population of over 700,000 (US Census Bureau 2010). This basin is impacted by commercial, agricultural, residential, and industrial discharges throughout the watershed (Page et al. 1992).

Substrates in the MS and MSC basins were highly variable ranging from predominately gravel, cobble and bedrock in the Clear Creek drainage, a sand-gravel mixture in the Mary's River drainage, to sand, silt, and gravel mixture in the MSC basin. Unstable shifting sand substrate (Figure 1) and claypan were recorded at a higher percentage in the MSC basin as compared to the other two basins. Shallow water depths, averaging 0.3 m, and ephemeral streams were predominant in the Clear Creek drainage (Figure 2), while the Mary's River drainage displayed stable substrates and a slightly deeper average water depth of 0.6 m.

Methods

During the 2009/2010 survey, freshwater mussel data were collected at 30 sites: 6 sites in Mary's River, 7 sites in Clear Creek, and 16 sites in the MSC basin (Figure 1). Locations of sampling sites are listed in Table 1 along with information regarding IDNR/IEPA sampling at the site. In most cases, mussel survey locations were the same as IDNR/IEPA sites.

Live mussels and shells were collected at each sample site to assess past and current freshwater mussel occurrences. Live mussels were surveyed by hand grabbing and visual detection (e.g. trails, siphons, exposed shell) when water conditions permitted. Efforts were made to cover all available habitat types present at a site including riffles, pools, slack water, and areas of differing substrates. A four-hour timed search method was implemented at each site. Live mussels were held in the stream until processing.

Following the timed search, all live mussels and shells were identified to species and recorded (Table 2). For each live individual, shell length (mm), gender, and an estimate of the number of growth rings recorded. Shell material was classified as recent dead (periostracum present, nacre pearly, and soft tissue may be present) or relict (periostracum eroded, nacre faded, shell chalky) based on condition of the best shell found. A species was considered extant at a site if it was represented by live or recently dead shell material (Szafoni 2001). The nomenclature employed in this report (Appendix 1) follows Turgeon et al. (1998) except for recent taxonomic changes to the gender ending of lilliput (*Toxolasma parvum*), which follows Williams et al. (2008). Voucher specimens were retained and deposited in the Illinois Natural History Survey Mollusk Collection. All non-vouchered live mussels were returned to the stream reach where they were collected.

Parameters recorded included extant and total species richness, presence of rare or listed species, and individuals collected, expressed as catch-per-unit-effort (CPUE) (Table 2). A population was considered to indicate recent recruitment if individuals less than 30 mm in length or with three or fewer growth rings were recorded. Finally, mussel resources were classified as Unique, Highly Valued, Moderate, Limited, or Restricted (Table 2) based on the above parameters (Table 3) and following criteria outlined in Table 4 (Szafoni, 2001).

Results

Species Richness

A total of 14 species of freshwater mussels were observed in the southern Mississippi basins, 10 of which were live (Table 2). Across all sites, the number of live species collected, the number of extant species collected (live + dead), and the total number of species collected (live + dead + relict) ranged from zero to six. The fragile papershell (*Leptodea fragilis*) had the most occurrences across sites sampled with live mussels present (four of nine sites; 44%; Figure 4). The giant floater (*Pyganodon grandis*), mapleleaf (*Quadrula quadrula*) and lilliput (*Toxolasma parvum*) were other commonly occurring species (Figure 4), occupying between 20% and 30% of these sites. Species richness varied greatly over the basin. Mary's River drainage displayed extant species richness (ESR) greater than four at 66% of the sites, while the Clear Creek drainage had only one site, Running Lake Ditch (site 27, ESR=4) with species counts above zero. Nearly 60% of the sites in the MSC and over 85% of the sites in the Clear Creek drainage did not support an extant mussel species assemblage. Site 23, Mary's River near Welge, had the greatest species richness with five live species.

Abundance and Recruitment

A total of 101 individuals were collected across 30 sites. The number of live specimens collected at a given site ranged from 0 to 52, with an average of nine mussels per site where live mussels were collected (9 of 30 sites; Table 2). A total of 84 collector-hours were spent sampling with an average of 2.8 mussels collected per hour at sites where mussels were present. Three sites (sites 10, 23, 27) yielded more than 10 individuals and one of those sites (site 23) yielded more than 50 live individuals. The most common species collected were mapleleaf (n=28), giant floater (n=24), white heelsplitter (*Lasmigona complanata*; n=15) and fragile papershell (n=12), which together comprised over 80% of the collections (Table 2). Nearly 80% of the live individuals were collected at two sites (site 23- Mary's River and site 10- Cahokia Canal).

Mussel abundance at individual sites ranged from none to moderate with CPUE ranging from 0 - 13 individuals/collector-hour (Table 2) Mary's River (site 23) was the only site in the basin to display a moderate CPUE at 13. All other sites had a limited CPUE. Mussel abundance was

ranked according to Table 2 and is shown in Figure 5.

Recruitment for each species was determined by the presence of individuals less than 30 mm or with 3 or fewer growth rings. Smaller (i.e., younger) mussels are harder to locate by hand grab methods and large sample sizes can be needed to accurately assess population reproduction. However, a small sample size can provide evidence of recruitment if it includes individuals that are small or possess few growth rings. Alternatively, a sample consisting of very large (for the species) individuals with numerous growth rings suggests a senescent population.

Recruitment at individual sites ranged from none observed to very high across the basin. Ninety percent of the sites (27 of 30) displayed no recruitment. Recruitment levels, referred to in Table 3 as Reproduction Factor, varied from one to five, with four sites exhibited high to very high recruitment (Table 3, Figure 5). Three sites in the Mary's River drainage, Cox Creek (site 20) and two mainstem sites (sites 19 and 23) had recruitment over 50% and 30-50%, respectively. Running Lake Ditch (site 27, 50% recruitment) was the only site in the Clear Creek drainage with recorded recruitment. The MSC system exhibited no observed recruitment during this survey.

Mussel Community Classification

Based on the data collected in the 2009/2010 basin surveys, many of the sites in the Mississippi South and South Central basins have Restricted or Limited mussel communities using the current MCI classification system (Table 4, Figure 5). Only four sites rank as Moderate mussel resources and one additional site (site 10) ranks as Limited. Three of the four Moderate mussel resource sites are in the Mary's River drainage and include two mainstem sites (sites 19 and 23) and Cox Creek (site 20). In the Clear Creek drainage, Running Lake Ditch (site 27) ranks as a Moderate mussel resource. One site in the MSC, Cahokia Canal (site 10), is listed as Limited; the 23 remaining sites in these basins are ranked as Restricted.

Noteworthy Finds

This survey collected 10 live species and 14 total species (live+ dead + relict). According to historical records, five species are known from the MSC basin, two species from the Clear Creek basin, and no historical data exists from the Mary's River drainage (Tiemann et al. 2007). To our knowledge, only 5 of the 30 sampled sites had previously been sampled for freshwater mussels. In the MSC basin, three new live species were recorded- mapleleaf, pink papershell (*Potamilus ohioensis*) and fragile papershell. New records for this basin also included two dead and three relict species: paper pondshell (*Utterbackia imbecillis*), pondhorn (*Uniomerus tetralasmus*), fatmucket (*Lampsilis siliquoidea*), cylindrical papershell (*Anodontoidea ferussacianus*), and threeridge (*Amblema plicata*). Three new species (two live, one dead shell) were detected in Running Lake Ditch (site 27, Clear Creek drainage) - giant floater, mapleleaf, and white heelsplitter. All mussels detected in the Mary's River drainage were new records as

no historical data were available from this basin (Table 2). Live records for this drainage included giant floater, white heelsplitter, mapleleaf, pondhorn, fragile papershell, lilliput, and yellow sandshell (*Lampsilis teres*).

Discussion

There is very limited historical mussel information from the Mississippi South and Mississippi South Central basins; nearly 85% of the sites sampled had no historical data available (Table 2), and no intensive survey for mussels has been completed in these drainages. The known historical data includes six species from five sites in these basins (Tiemann et al. 2007). Our surveys documented the existence of 14 species in the MS/MSC basins, and 10 of these species were represented by live individuals. All historically known species were found live during the 2009/2010 surveys at one or more sites except paper pondshell, which was represented at three sites by dead or relict shells (Table 2). Five additional live species were collected during this survey including mapleleaf, pondhorn, white heelsplitter, yellow sandshell, and pink papershell. These species are widespread and fairly common throughout the state. New species records were collected at 16 of the 30 sites, including 5 of the 6 sites in the Mary's River Basin.

Recruitment

Only four sites in the MS exhibited high to very high recruitment; these included three Mary's River drainage sites (two mainstem and Cox Creek) and Running Lake Ditch in the Clear Creek drainage. This finding suggests that the mussel communities of the Mary's River are viable and self-maintaining at this time. Data collected during this survey indicate that very recent recruitment may not be occurring at any site in the MSC basin or in 85% of the Clear Creek drainage. Sampling methods to target juvenile mussels would be necessary to better assess the reproductive status of these populations.

Mussel community of the Mississippi South/South Central basin

Based on limited historical information it is uncertain if the mussel communities of the MSC and MS basins have changed over time. In the MSC basin, only 7 out of 17 sites had an extant population greater than 0 and 6 of those sites contained only 1 to 2 extant species. Relict shells were recorded at 15 of the 17 sites in this basin and the majority of 8 species detected were recorded as dead or relict shell. From this survey, it would appear that the majority of the MSC region does not support extant mussel communities. Due to minimal historical data, it cannot be determined whether the MSC basin lacks mussel diversity because streams lack suitable habitat or if it has suffered species extirpations. The presence of relict shell at a large percentage of sites (88%) points towards the extirpation of species in this drainage.

Urbanization and industrialization have impacted many streams in this basin, and have led to alterations in stream side vegetation cover, loss of instream habitat, and changes in stream depth and velocity. Other factors that may be affecting stream ecosystems in this area include elevated concentrations of fecal coliform, phosphorus, iron, manganese, and aquatic algae (IEPA 2010). The combination of these factors may be having a detrimental effect on mussel communities in this basin.

In the Clear Creek drainage, 85% of the streams lacked mussel occurrence. Live, dead, and relict shells were recorded from only one stream in the drainage, Running Lake Ditch, which ranked as a Moderate mussel resource. Nearly all of the streams in this drainage are considered full support for aquatic life based on biological, physiochemical, physical habitat, and toxicity data collected by the IEPA (IEPA 2010). Although able to support aquatic life based on these standards, many of the streams sampled in this drainage are considered ephemeral and may lack the water availability needed to support a mussel community.

In the Mary's River drainage, live specimens, or dead and relict shells were collected at five of six sites sampled. Piney Branch, a bedrock laden ephemeral creek located in Piney Branch Ravine, was the only site in the drainage with no mussel presence detected. Three of the streams in the Mary's River drainage appear to support a moderate mussel fauna and high to very high recruitment at this time. This suggests that the mussel communities of the Mary's River drainage appear to be viable and self-maintaining. Located in close proximity to the diverse mussel communities of the Kaskaskia River basin, it seems reasonable to assume that the Mary's River drainage could be capable of supporting a biologically significant freshwater mussel fauna in the future.

Literature Cited

Bogan, A.E. 1993. Freshwater bivalve extinctions (Mollusca: Unionoida): a search for causes. *American Zoologist* 33(6):599-609.

Illinois Endangered Species Protection Board. 2011. Checklist of endangered and threatened animals and plants of Illinois. http://dnr.state.il.us/ESPB/pdf/2011_Checklist.pdf

Illinois Environmental Protection Agency. 2010. Illinois Integrated Water Quality Report and Section 303(d) List. <http://www.epa.state.il.us/water/tmdl/303d-list.html>

Illinois Department of Agriculture. 2000. Land Cover of Illinois Statistical Summary 1999-2000. <http://www.agr.state.il.us/gis/stats/landcover99-00.html>

Page, L.M., K.S. Cummings, C.A. Mayer, S.L. Post, and M.E. Retzer. 1992. Biologically significant Illinois streams, an evaluation of the streams of Illinois based on aquatic biodiversity. Technical Report. Illinois Department of Conservation and Illinois Department of Energy and Natural Resources, Springfield, Illinois. 498 pp.

Schwegman, J.E. 1973. Comprehensive plan for the Illinois nature preserves system. Part 2. The natural divisions of Illinois. Illinois Nature Preserves Commission, Springfield, Illinois.

Strayer, D.L., J.A. Downing, W.R. Haag, T.L. King, J.B. Layzer, T.J. Newton, and S.J. Nichols. 2004. Changing perspective on pearlymussels, North America's most imperiled animals. *BioScience* 54(5):429-439.

Szafoni, R. E. 2001. Protocol for integrating freshwater mussel surveys into IDNR / IEPA stream basin surveys. Version 2.0. IDNR/ORC/Natural Heritage, Charleston, IL. 5pp.

Tiemann, J.S., K.S. Cummings, C.A. Mayer. 2007. Updates to the distributional checklist and status of Illinois freshwater mussels (Mollusca: Unionacea). *Transactions of the Illinois State Academy of Science* 100 (1):107-123.

Turgeon, D.D., A.E. Bogan, E.V. Coan, F.G. Hochberg, W.G. Lyons, P.M. Mikkelsen, J.F. Quinn, Jr., C.F.E. Roper, G. Rosenberg, B. Roth, A. Scheltema, M.J. Sweeney, F.G. Thompson, M. Vecchione, and J.D. Williams. 1998. Common and scientific names of aquatic invertebrates from the United States and Canada: Mollusks. 2nd Edition. American Fisheries Society, Special Publication 26:ix-526.

U.S. Census Bureau; 2010 Census Fact finder. <http://factfinder2.census.gov>

Williams, J.D., M.L. Warren, Jr., K.S. Cummings, J.L. Harris, and R.J. Neves. 1993. Conservation status of freshwater mussels of the United States and Canada. *Fisheries* 18(9):6-22.

Williams, J.D., A.E. Bogan, and J.T. Garner. 2008. The freshwater mussels of Alabama and the Mobile Basin of Georgia, Mississippi, and Tennessee. University of Alabama Press, Tuscaloosa, Alabama. 908 pp.

Table 1. 2009-2010 Mississippi South/South Central Intensive Basin Survey. Types of samples include MU-mussel sampling, BE-boat electrofishing, ES-electric fish seine, FF-fish flesh contaminate, H-habitat, M-macroinvertebrate, S-sediment, W-water chemistry.

| Site Number | IEPA Code | Stream | Types of Sample | County | Location | Watershed Area (km ²) |
|----------------------------------|-----------|--------------------|------------------|-----------|---|-----------------------------------|
| Mississippi South Central | | | | | | |
| 1 | JQ-04 | Cahokia Creek | MU,ES,M,H,S,W | Macoupin | Rt 138, 2 mi SE Benld | 112.76 |
| 2 | JQE-01 | Sherry Creek | MU,ES,M,H,S,W | Madison | Possum Hill Rd, E of Holiday Shores | 71.69 |
| 3 | JV-02 | Piasa Creek | MU,ES,M,H,S,W | Madison | Airport Road, NW of Alton | 155.77 |
| 4 | JVA-01 | Mill Creek | MU,SH,M,H,S,W | Jersey | Lockhaven Rd, 3 mi E Elsah | 37.76 |
| 5 | JRB-01 | W Fk Wood River | MU,SH,M,H,S,W | Madison | Wood Station Rd, 1 mi N Alton | 100.40 |
| 6 | JRA-02 | E Fk Wood River | MU,SH,M,H,S,W | Madison | Culp Lane, NW edge Bethalto | 113.84 |
| 7 | JR-02 | Wood River | MU,ES,M,H,S,W | Madison | Rt 3, at Milton Rd East Alton | 312.01 |
| 8 | JQA-01 | Indian Creek | MU | Madison | Rt 143, 2 Mi E Wood River | 104.76 |
| 9 | JQ-05 | Cahokia Creek | MU,ES,FF,M,H,S,W | Madison | Rt 143, NW Edwardsville | 544.78 |
| 10 | JN-02 | Cahokia Canal | MU,ES,M,H,S,W | Madison | N Br, Sand Prairie Ln Horseshoe Lk | 196.02 |
| 11 | JNA-01 | Canteen Creek | MU,ES,M,H,S,W | Madison | S Br, Sand Prairie Ln Horseshoe Lk | 68.61 |
| 12 | JMAC-02 | Harding Ditch | MU,ES,M,H,S,W | St. Clair | Lake Dr, Frank Holten St Pk | 84.95 |
| 13 | JMAA-03 | Prairie DuPont | MU,ES,M,H,S,W | St. Clair | Zingg Road, 1.5 mi N Millstadt | 39.44 |
| 14 | JI-01 | Carr Creek | MU,ES,M,H,S,W | Monroe | Valmeyer Rd, W of Rt 3, Columbia | 16.91 |
| 15 | JH-03 | Fountain Creek | MU | Monroe | HH Rd Br, UPS Andys Run | 98.09 |
| 16 | JH-04 | Fountain Creek | MU,ES,M,H,S,W | Monroe | Rt. 156 Br, 2mi W Waterloo | 65.42 |
| 17 | JD-02 | Maeystown Creek | MU,ES,M,H,S,W | Monroe | Bluff Rd Br, NE Chalfin Br. | 25.81 |
| Mary's River | | | | | | |
| 18 | IJK-27 | Maxwell Creek | MU,ES,H,M,S,W | Randolph | 2 Mi SE Sparta on Pierce Lane | 23.48 |
| 19 | II-91 | Marys River | MU,ES,H,M,W | Randolph | Eden Rd, 1.5 Mi N Steeleville | 126.35 |
| 20 | IIH-36 | Cox Creek | MU,ES,H,M,W | Randolph | Chester St, 0.5 Mi S Steeleville | 261.28 |
| 21 | IIC-39 | Little Marys River | MU,ES,H,M,W | Randolph | 3 Mi NE Chester At Covered Br | 178.13 |
| 22 | IIBB-01 | Piney Branch | MU,ES,H,M,S,W | Randolph | Piney Creek Road 2.3 Mi SW of Shiloh Hi | 13.66 |
| 23 | II-03 | Marys River | MU,BE,H,M,W | Randolph | 0.5 Mi E of Welge | 288.88 |
| Clear Creek | | | | | | |
| 24 | ICE-02 | Hutchins Creek | MU,ES,H,M,S,W | Union | Beech Grove Rd, 5 mi. SW Alto Pass | 52.78 |
| 25 | IC-02 | Clear Creek | MU,ES,H,M,W | Union | 1 Mi SSE of Bald Knob | 46.28 |
| 26 | ICDB-01 | Green Creek | MU,ES,H,M,S,W | Union | Next To Rt 146 Br, 2.9 Mi W Jonesboro | 23.54 |
| 27 | ICIA-02 | Running Lake Ditch | MU,ES,H,M,S,W | Union | Ware-Wolf Lake Rd, 2 Mi W of Wolf Lake | 61.52 |
| 28 | ICD-01 | Dutch Creek | MU,ES,H,M,S,W | Union | Farm Rd, 2 Mi W Jonesboro | 50.93 |
| 29 | IB-03 | Sexton Creek | MU,ES,H,M,W | Alexander | 2.8 Mi NE of Gale | 32.15 |
| 30 | IBAA-01 | Sammons Creek | MU,ES,H,M,S,W | Alexander | Bean Ridge Rd 1.7 Mi NE of Thebes | 8.63 |

Table 3. Mussel Community Index (MCI) parameters and scores.

| | | | |
|---|------------------------|---------------------------------|------------------------------|
| Extant species in sample | Species Richness | Catch per Unit Effort (CPUE) | Abundance (AB) Factor |
| 0 | 1 | 0 | 0 |
| 1-3 | 2 | 1-10 | 2 |
| 4-6 | 3 | >10-30 | 3 |
| 7-9 | 4 | >30-60 | 4 |
| 10+ | 5 | >60 | 5 |
| % live species with recent recruitment | Reproduction Factor | # of Intolerant species | Intolerant species Factor |
| 0 | 1 | 0 | 1 |
| 1-30 | 3 | 1 | 3 |
| >30-50 | 4 | 2+ | 5 |
| >50 | 5 | | |

Table 4. Freshwater mussel resource categories based on species richness, abundance, and population structure. MCI = Mussel Community Index Score

| | |
|--|--|
| Unique Resource MCI \geq 16 | Very high species richness (10 + species) &/or abundance (CPUE > 80); intolerant species typically present; recruitment noted for most species |
| Highly Valued Resource MCI = 12- 15 | High species richness (7-9 species) &/or abundance (CPUE 51-80); intolerant species likely present; recruitment noted for several species |
| Moderate Resource MCI = 8 - 11 | Moderate species richness (4-6 species) &/or abundance (CPUE 11-50) typical for stream of given location and order; intolerant species likely not present; recruitment noted for a few species |
| Limited Resource MCI = 5 - 7 | Low species richness (1-3 species) &/or abundance (CPUE 1-10); lack of intolerant species; no evidence of recent recruitment (all individuals old or large for the species) |
| Restricted Resource MCI = 0 - 4 | No live mussels present; only weathered dead, sub-fossil, or no shell material found |




Figure 1. Sites sampled in the Mississippi South and Mississippi South Central basins during 2009- 2010. Site codes referenced in Table 1.



Figure 2. East Fork Wood River in the Mississippi South Central basin. Substrate predominately shifting unstable sand, large sand pile can be seen in the upper right hand corner of the picture.



Figure 3. Sammons Creek in the Mississippi South basin. Substrates predominately small to medium gravel mix, water levels reduced to intermittent pools.




Figure 4. Number of sites where a species was collected live compared to the number of sites sampled with live specimens (9 total sites).




Figure 5. Comparison of Mussel Community Index (MCI) and MCI component scores for Mississippi South and Mississippi South Central River basin sites based on factor values from Table 3.

Appendix 1. Scientific and common names of species.

| Scientific name | Common name |
|-----------------------------------|--------------------|
| Subfamily Anodontinae | |
| <i>Anodonta suborbiculata</i> | flat floater |
| <i>Anodontoides ferussacianus</i> | |
| <i>Lasmigona complanata</i> | white heelsplitter |
| <i>Pyganodon grandis</i> | giant floater |
| <i>Utterbackia imbecillis</i> | paper pondshell |
| Subfamily Ambleminae | |
| <i>Amblema plicata</i> | threeridge |
| <i>Quadrula quadrula</i> | mapleleaf |
| <i>Unio merus tetralasmus</i> | pondhorn |
| Subfamily Lampsilinae | |
| <i>Lampsilis siliquoidea</i> | fatmucket |
| <i>Lampsilis teres</i> | yellow sandshell |
| <i>Leptodea fragilis</i> | fragile papershell |
| <i>Ligumia subrostrata</i> | pondmussel |
| <i>Potamilus ohioensis</i> | pink papershell |
| <i>Toxolasma parvum</i> | lilliput |